

Creating strong bonds through a full range of leading vaccines



Nobivac: Rabies

 Dog

 Cat

 Core

 Non Core

 Inactivated

 DOI

A LEADING VACCINE IN THE FIGHT AGAINST RABIES

- Proven 3-Year DOI
- Used worldwide as an approved choice for many national and regional rabies control programs
- Used in the highly successful Serengeti rabies control program leading to a major reduction in both canine and human rabies cases
- Can be given together with many other Nobivac vaccines

Nobivac: Puppy DP






 Dog

 Core

 Live

TO SUPPORT A SOCIAL START

- Can be used from 4-6 weeks of age
- Provides early CDV and CPV immunity
- Breaks through maternally derived antibodies earlier
- Reduces the disease risk for early socialization programmes – makes early social interactions with other dogs safer

ICONS	
	Live Vaccine
	Core Vaccination
	Inactivated Vaccine
	Non Core Vaccination
	DOI: Duration Of Immunity





Nobivac:DHPPi



PROVIDES WHAT EVERY DOG NEEDS

- Reliable immunity against distemper, infectious hepatitis, canine parvovirus disease and respiratory disease caused by canine adenovirus type 2 and canine parainfluenza virus*
- Logical antigen combinations mean fewer injections to cover basic vaccine needs
- Fast onset of immunity within 7 days for D, H and P core components
- Can be used in pregnant and lactating bitches if required
- Proven clinical protection against CPV-2, 2a, 2b and 2c strains & prevention of viral shedding^{1, 2, 3}
- 3-Year duration of immunity for DH and P core components

Nobivac:Lepto



REDUCES RENAL SHEDDING

- For active immunity against leptospirosis caused by *Leptospira canicola* and *Leptospira icterohaemorrhagiae*
- First leptospirosis vaccine with an authorized claim to reduce renal shedding; thus reducing the risk of zoonotic spread
- Non-adjuvanted vaccine
- 1-year DOI**

¹ Hoskins J.D., Taylor H.W. and Gourley K.R., (1995), Challenge trial of a new attenuated canine parvovirus vaccine, *J.Vet.Int.Med.*, **9**, 197.

² Larson L.J. and Schultz R.D., (1997) Comparison of selected canine vaccines for their ability to induce protective immunity against canine parvovirus infection, *American Journal of Veterinary Research*, **58**, 360-363.

³ Spibey N., Greenwood N.M., Sutton D., Chalmers W.S.K. and Tarpey I., (2008), Canine parvovirus type 2 protects against virulent challenge with type 2c virus, *Veterinary Microbiology*, **128**, 48-55

Nobivac Tricat Trio

Feline Vaccination Protocols



THE CORE VACCINE FOR CATS

- Immunizes against feline calicivirus (FCV), feline herpes (or rhinotracheitis) virus type 1 (FHV) and feline panleucopenia virus (FPLV)
- Highly efficacious live attenuated strains
- Non-adjuvanted vaccine
- 3-Year DOI for feline panleucopenia component
- Contains the broadly cross-protective F9 FCV component. Shown to cross neutralize a wide range of FCV field isolates⁴

Core vaccination for cats: FHV, FCV and FPLV

+ Rabies



	Tricat Trio*
6w	
8-9w	X
12w	X
16-20w	
1 st year	X
2 nd year	X
3 rd year	X
4 th year	X
5 th year	X
6 th year	X
7 th year	X
8 th year	X

	Rabies**
6w	
8-9w	
12w	X
16-20w	
1 st year	(X)
2 nd year	(X)
3 rd year	X
4 th year	(X)
5 th year	(X)
6 th year	X
7 th year	(X)
8 th year	(X)

*Vaccine has 3-year claim for FPLV but FHV and FCV needs boosting annually
**Rabies annual booster depends on the country's regulatory requirements and not necessary according to our SPC (it has 3-year claim).



🐾

Canine Vaccination Protocols

Core vaccination for dogs: DH and P		+ Lepto		+ Rabies	
	<div>●</div> Puppy DP*	<div>●</div> DHPPI**		<div>●</div> Lepto	<div>●</div> Rabies***
6w	(X)		6w		
8-9w		X	8-9w	X	
12w		X	12w	X	X
16-20w			16-20w		
1 st year		X	1 st year	X	(X)
2 nd year		X	2 nd year	X	(X)
3 rd year		X	3 rd year	X	X
4 th year		X	4 th year	X	(X)
5 th year		X	5 th year	X	(X)
6 th year		X	6 th year	X	X
7 th year		X	7 th year	X	(X)
8 th year		X	8 th year	X	(X)

*Nobivac solvent must be used to reconstitute Nobivac Puppy DP and can be used to reconstitute Nobivac DHPPI.

**Vaccines have 3-year claim for D,H and P but Pi needs boosting annually.

***Rabies annual booster depends on the country's regulatory requirements and not necessary according to our SPC (it has 3-year claim).





Protecting strong bonds

The Nobivac small animal vaccine range comprises vaccines and services that reflect MSD Animal Health long experience and leadership in small animal vaccine development. A strong research commitment to small animal vaccines ensure that the Nobivac vaccines are of the highest quality, trusted to be effective and safe.

